

University of Groningen

## Immobilization of rhodium complexes at thiolate monolayers on gold surfaces

Belser, T; Stöhr, Meike; Pfaltz, A

*Published in:*  
Journal of the American Chemical Society

*DOI:*  
[10.1021/ja0500714](https://doi.org/10.1021/ja0500714)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2005

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Belser, T., Stöhr, M., & Pfaltz, A. (2005). Immobilization of rhodium complexes at thiolate monolayers on gold surfaces: Catalytic and structural studies. *Journal of the American Chemical Society*, 127(24), 8720-8731. <https://doi.org/10.1021/ja0500714>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Immobilization of Rhodium Complexes at Thiolate Monolayers on Gold Surfaces: Catalytic and Structural Studies


Thomas Belser<sup>†</sup>, Meike Stöhr<sup>‡,\*</sup> and Andreas Pfaltz<sup>†,\*</sup>

Contribution from the Department of Chemistry,<sup>†</sup> St. Johannis-Ring 19 and the Department of Physics,<sup>‡</sup> Klingelbergstrasse 82, University of Basel, 4056 Basel (Switzerland).

## SUPPORTING INFORMATION

**General Methods:** Reactions with air- or moisture-sensitive compounds were performed under argon using standard Schlenk techniques or under purified N<sub>2</sub> in a MBraun glovebox. Glasware was oven dried and flame dried prior to use. All chemicals were purchased from Fluka or Aldrich. CH<sub>2</sub>Cl<sub>2</sub> was dried over CaH<sub>2</sub> and THF over Na/K and freshly distilled under a stream of nitrogen prior to use. Optical rotations were measured in a Perkin Elmer Polarimeter 341, sodium lamp, 1 dm cuvette length, *c* in g/100 mL. <sup>1</sup>H-, <sup>13</sup>C-, <sup>19</sup>F- and <sup>31</sup>P-NMR spectra were recorded in CD<sub>2</sub>Cl<sub>2</sub> or CDCl<sub>3</sub> on a Bruker Avance 400 MHz or 500 MHz spectrometer and coupling constants are reported in Hz. Chemical shifts are given in ppm relative to TMS. IR spectra: 1600 Perkin Elmer Series FT-IR spectrometer; FAB mass spectra: Finnigan MAT 312; EI mass spectra: VG 70 SE. Elemental analysis were carried out on a Leco CHN-900 and Leco RO-478. Chromatographic purifications were performed by flash chromatography using silica gel (Merck 0.040 - 0.063 mm). Yields for final products refer to isolated products and are the average of three runs. GC analyses: Carlo Erba HRGC Mega2 Series MFC 800 (column: Restek Rtx-1701; 0.25 mm, 30 m, 60 kPa He). HPLC analyses: Shimadzu VP-system (column: Daicel OD-H; 4.6 × 250 mm).

### Synthesis of 11-(10'-carboxy-decyldisulfanyl)undecanoic acid



Sodium hydroxide (366 mg, 9.16 mmol), potassium iodide (50.0 mg, 0.30 mmol) and iodine (1.16 g, 4.58 mmol) were added to a solution of 11-mercaptoundecanoic acid (2.00 g, 9.16 mmol) in methanol (50 mL) and the solution was stirred for 30 min. The brown reaction mixture was decolorized with a saturated sodium sulfite solution, the solvents concentrated under reduced pressure, the precipitate dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and the resulting solution washed with a HCl solution (1 M, 50 mL) and water (50 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced

pressure. Purification of the crude material by crystallization (hexanes/AcOEt) gave 63% yield of the desired 11-(10'-carboxy-decyl)disulfanylundecanoic acid (1.25 g).

m.p.: 68 - 70 °C.  $R_f$  = 0.30 (hexanes/AcOEt = 4:1).

$^1\text{H}$ -NMR (400.1 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 1.28 ( $s_b$ , 20 H,  $\text{CH}_2$ ), 1.35 ( $m_c$ , 4 H,  $\text{CH}_2$ ), 1.64 ( $m_c$ ,  $J$  = 7.4 Hz, 8 H,  $\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$  +  $\text{SCH}_2\text{CH}_2$ ), 2.34 (t,  $J$  = 7.4 Hz, 4 H,  $\text{CH}_2\text{CO}_2\text{H}$ ), 2.68 (t,  $J$  = 7.4 Hz, 4 H,  $\text{SCH}_2$ ), 10.98 ( $s_b$ , 2 H,  $\text{CO}_2\text{H}$ ).


$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 25.0 ( $\text{CH}_2$ ), 28.9 ( $\text{CH}_2$ ), 29.4 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 29.6 ( $\text{CH}_2$ ), 29.7 ( $\text{CH}_2$ ), 29.8 ( $\text{CH}_2$ ), 34.4 ( $\text{CH}_2$ ), 39.6 ( $\text{SCH}_2$ ), 180.7 ( $\text{CO}_2\text{H}$ ).

IR (KBr):  $\nu$  ( $\text{cm}^{-1}$ ) = 2918s, 2849s, 1696s, 1471m, 1427m, 1289m, 1261m, 1231m, 1206m, 944m.

MS (FAB):  $m/z$  (rel int %) = 434 ( $\text{M}^+$ , 30), 417 (20), 344 (34), 328 (15), 232 (13), 171 (15), 127 (14), 111 (27), 101 (25), 87 (57), 81 (32), 69 (49), 55 (100), 43 (47).

Elemental analysis for  $\text{C}_{22}\text{H}_{42}\text{O}_4\text{S}_2$  calc.: C 60.79%, H 9.74%, O 14.72%; found: C 60.84%, H 9.68%, O 14.74%.

### Synthesis of (3*R*,4*R*)-3,4-bis(diphenylphosphorothioyl)pyrrolidine



Sulfur (0.27 g, 8.40 mmol) was added to a solution of PYRPHOS (1.85 g, 4.20 mmol) in freshly distilled toluene (50 mL) and the reaction mixture was stirred for 12 hours at 23 °C. The resulting solution was concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with diethyl ether, gave 95% yield of the desired compound (2.01 g).

m.p.: 212 - 214 °C.  $[\alpha]_D^{20}$  = +12.3 ( $c$  = 0.80,  $\text{CHCl}_3$ ).  $R_f$  = 0.50 (diethyl ether).

$^1\text{H}$ -NMR (400.1 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 2.59 ( $s_b$ , 1 H,  $\text{NH}$ ), 3.12 ( $m_c$ , 2 H,  $\text{NCH}_2$ ), 3.27 ( $m_c$ , 2 H,  $\text{NCH}_2$ ), 4.01 ( $m_c$ , 2 H,  $\text{PCH}$ ), 7.00 ( $m_c$ , 4 H,  $\text{CH}_{\text{Ph}}$ ), 7.17 ( $m_c$ , 2 H,  $\text{CH}_{\text{Ph}}$ ), 7.42 ( $m_c$ , 6 H,  $\text{CH}_{\text{Ph}}$ ), 7.60 ( $m_c$ , 4 H,  $\text{CH}_{\text{Ph}}$ ), 7.88 ( $m_c$ , 4 H,  $\text{CH}_{\text{Ph}}$ ).

$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 39.3 - 40.1 (dd,  $\text{PCH}$ ), 53.1 ( $\text{NCH}_2$ ), 128.3 - 128.8 (m,  $\text{CH}_{\text{Ph-meta}}$ ), 130.0 - 132.9 (m,  $\text{C}_{\text{Ph-ipso}}$ ), 130.9 - 131.6 (m,  $\text{CH}_{\text{Ph-ortho and para}}$ ).

$^{31}\text{P}\{^1\text{H}\}$ -NMR (162.0 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 51.2 (s).


IR (KBr):  $\nu$  ( $\text{cm}^{-1}$ ) = 3300m, 3048m, 2898m, 1479m, 1435s, 1311m, 1233w, 1158m, 1099s, 998m, 861s, 745s, 718s, 693s, 646s, 605m, 566m, 521s, 503s, 489s.

MS (FAB):  $m/z$  (rel int %) = 504 ( $[\text{M} + \text{H}]^+$ , 30), 286 (9), 217 (15), 185 (6), 77 (5), 68 (100), 39 (5).

Elemental analysis for  $\text{C}_{28}\text{H}_{27}\text{NP}_2\text{S}_2$  calc.: C 66.78%, H 5.40%, N 2.78%; found: C 66.72%, H 5.32%, N 2.75%.

## SYNTHESIS OF THE SPACERS FOR THE GOLD COLLOIDS

### Synthesis of 1-(2'-bromo-ethyl)adamantane



*N*-Bromosuccinimide (987 mg, 5.55 mmol) and triphenylphosphine (1.46 g, 5.55 mmol) were added to a solution of 2-adamant-1-yl-ethanol (1.00 g, 5.55 mmol) in freshly distilled benzene (20 mL) and the reaction mixture was stirred for 12 hours at 23 °C. The resulting solution was washed with a saturated NaCl solution (20 mL) and water (20 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with *n*-pentane, gave 92% yield of 1-(2'-bromo-ethyl)adamantane (1.24 g).

m.p.: 66 - 68 °C. *R*<sub>f</sub> = 0.78 (*n*-pentane).

<sup>1</sup>H-NMR (400.1 MHz, CDCl<sub>3</sub>, 295 K): δ = 1.51 (m<sub>c</sub>, *J* = 2.5 Hz, 6 H, CCH<sub>2-Ad</sub>), 1.61 - 1.70 (m, 6 H, CH<sub>2-Ad</sub>), 1.72 (m<sub>c</sub>, 2 H, BrCH<sub>2</sub>CH<sub>2</sub>), 1.96 (m<sub>c</sub>, 3 H, CH<sub>Ad</sub>), 3.40 (m<sub>c</sub>, 2 H, BrCH<sub>2</sub>).


<sup>13</sup>C{<sup>1</sup>H}-NMR (100.6 MHz, CDCl<sub>3</sub>, 295 K): δ = 28.6 (CH<sub>Ad</sub>), 29.2 (BrCH<sub>2</sub>), 34.1 (C<sub>Ad</sub>), 37.1 (CH<sub>2-Ad</sub>), 42.2 (CCH<sub>2-Ad</sub>), 48.2 (BrCH<sub>2</sub>CH<sub>2</sub>).

IR (KBr): ν (cm<sup>-1</sup>) = 2896s, 2843s, 2652w, 1452s, 1442m, 1344m, 1329m, 1312m, 1256m, 1215m, 1094m, 973m, 810m, 746w, 719m, 658m, 560s.

MS (EI): *m/z* (rel int %) = 242 (M<sup>+</sup>, 1, <sup>79</sup>Br), 135 (100), 107 (6), 93 (13), 79 (15), 67 (6), 55 (3), 41 (6).

Elemental analysis for C<sub>12</sub>H<sub>19</sub>Br calc.: C 59.27%, H 7.88%; found: C 59.22%, H 7.79%.

### Synthesis of 1-(hex-5'-enyl)adamantane



3-Butenylmagnesium bromide solution (0.5 M in THF, 94 mL, 47.0 mmol) was added tropwise at -78 °C to a solution of 1-(2'-bromo-ethyl)adamantane (5.00 g, 20.6 mmol) in freshly distilled THF (30 mL). After 10 min, a lithium tetrachlorocuprate(II) solution (0.1 M in THF, 2.10 mL, 0.21 mmol) was added and the reaction mixture was stirred for 2 hours at -78 °C and slowly warmed up to 23 °C over 12 hours. The resulting solution was washed with a saturated NH<sub>4</sub>Cl solution (100 mL) and water (100 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with *n*-pentane, gave 92% yield of 1-(hex-5'-enyl)adamantane (4.13 g).

*R*<sub>f</sub> = 0.86 (hexanes).

<sup>1</sup>H-NMR (400.1 MHz, CDCl<sub>3</sub>, 295 K): δ = 1.03 (t, *J* = 7.1 Hz, 2 H, CCH<sub>2</sub>), 1.24 (quin, *J* = 7.1 Hz, 2 H, CCH<sub>2</sub>CH<sub>2</sub>), 1.33 (quin, *J* = 7.1 Hz, 2 H, CCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.45 (m<sub>c</sub>, *J* = 2.2 Hz, 6 H, CCH<sub>2-Ad</sub>), 1.66

(m<sub>c</sub>, 6 H, CH<sub>2-Ad</sub>), 1.93 (m<sub>c</sub>, 3 H, CH<sub>Ad</sub>), 2.05 (m<sub>c</sub>, 2 H, CCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>) 4.96 (m<sub>c</sub>, 2 H, CH<sub>2-vinyl</sub>), 5.81 (m<sub>c</sub>, 1 H, CH<sub>vinyl</sub>).


<sup>13</sup>C{<sup>1</sup>H}-NMR (100.6 MHz, CDCl<sub>3</sub>, 295 K): δ = 22.0 (CH<sub>2</sub>), 28.9 (CH<sub>Ad</sub>), 30.0 (CH<sub>2</sub>), 32.4 (C<sub>Ad</sub>), 34.0 (CH<sub>2</sub>), 37.4 (CH<sub>2-Ad</sub>), 42.7 (CCH<sub>2-Ad</sub>), 44.7 (CH<sub>2</sub>), 114.2 (CH<sub>2-vinyl</sub>), 139.4 (CH<sub>vinyl</sub>).

IR (NaCl): ν (cm<sup>-1</sup>) = 3076w, 2898s, 2845s, 2656w, 1820w, 1640m, 1450m, 1098w, 991m, 908m.

MS (EI): *m/z* (rel int %) = 218 (M<sup>+</sup>, 3), 135 (100), 107 (4), 93 (8), 79 (9), 67 (5), 55 (4), 41 (5).

Elemental analysis for C<sub>16</sub>H<sub>26</sub> calc.: C 88.00%, H 12.00%; found: C 88.07%, H 12.01%.

### Synthesis of thioacetic acid (*S*)-6-(adamant-1'-yl)hexyl ester



Thioacetic acid (4.21 mL, 58.9 mmol) was added to a solution of 1-(hex-5'-enyl)adamantane (5.00 g, 22.9 mmol) in freshly distilled THF (250 mL) and cooled to 20 °C with a cryostate (cooling bath: H<sub>2</sub>O/EtOH = 80:20). The solution was irradiated during 7 hours with a 300 W lamp (Osram Ultra-Vitalux 300 W). During that time was added 2,2'-azobisisobutyronitrile (900 mg, 5.48 mmol) in freshly distilled THF (20 mL) with a syringe pump. Irradiation was continued for another 4 hours. The solvent was concentrated under reduced pressure and the residue dissolved in dichloromethane (100 mL). The resulting solution was washed with a saturated Na<sub>2</sub>CO<sub>3</sub> solution (100 mL) and water (100 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with 5% diethyl ether in *n*-pentane, gave 82% yield of thioacetic acid (*S*)-6-(adamant-1'-yl)hexyl ester (5.50 g).

R<sub>f</sub> = 0.67 (*n*-pentane/diethyl ether = 20:1).

<sup>1</sup>H-NMR (400.1 MHz, CDCl<sub>3</sub>, 295 K): δ = 1.00 (t, *J* = 7.3 Hz, 2 H, CCH<sub>2</sub>), 1.22 (m<sub>c</sub>, 4 H, CH<sub>2</sub>), 1.35 (quin, *J* = 7.3 Hz, 2 H, CCH<sub>2</sub>CH<sub>2</sub>), 1.43 (m<sub>c</sub>, *J* = 2.3 Hz, 6 H, CCH<sub>2-Ad</sub>), 1.55 (quin, *J* = 7.3 Hz, 2 H, CH<sub>2</sub>CH<sub>2</sub>S), 1.64 (m<sub>c</sub>, 6 H, CH<sub>2-Ad</sub>), 1.92 (m<sub>c</sub>, 3 H, CH<sub>Ad</sub>), 2.31 (s, 3 H, SCOCH<sub>3</sub>), 2.85 (t, *J* = 7.3 Hz, 2 H, CH<sub>2</sub>S).


<sup>13</sup>C{<sup>1</sup>H}-NMR (100.6 MHz, CDCl<sub>3</sub>, 295 K): δ = 22.3 (CH<sub>2</sub>), 28.8 (CH<sub>Ad</sub>), 29.0 (CH<sub>2</sub>), 29.3 (CH<sub>2</sub>), 29.6 (CH<sub>2</sub>), 30.2 (CH<sub>2</sub>), 30.7 (CH<sub>3</sub>), 32.3 (C<sub>Ad</sub>), 37.4 (CH<sub>2-Ad</sub>), 42.6 (CCH<sub>2-Ad</sub>), 44.7 (CH<sub>2</sub>), 196.1 (SCO).

IR (NaCl): ν (cm<sup>-1</sup>) = 2902s, 2847s, 2661w, 1694s, 1450m, 1352m, 1132m, 954m, 864w, 624w.

MS (EI): *m/z* (rel int %) = 294 (M<sup>+</sup>, 4), 251 (5), 135 (100), 107 (5), 93 (9), 79 (10), 43 (15).

Elemental analysis for C<sub>18</sub>H<sub>30</sub>OS calc.: C 73.41%, H 10.27%, O 5.43%; found: C 73.21%, H 10.14%, O 5.57%.

## Synthesis of 6-(adamant-1'-yl)hexanethiol



Conc. HCl (37%, 425 mL, 5.18 mmol) was added to a solution of thioacetic acid (*S*)-6-(adamant-1'-yl)-hexyl ester (150 mg, 509  $\mu$ mol) in methanol (5 mL) and the reaction mixture was stirred for 10 hours at 65 °C. The solvent was concentrated under reduced pressure and the residue dissolved in diethyl ether (20 mL). The resulting solution was washed with a HCl solution (1 M, 20 mL) and water (20 mL). The organic phase was dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with *n*-pentane, gave 86% yield of 6-(adamant-1'-yl)-hexanethiol (0.11 g).

$R_f = 0.65$  (*n*-pentane).

$^1\text{H}$ -NMR (400.1 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta = 1.01$  (t,  $J = 7.4$  Hz, 2 H,  $\text{CCH}_2$ ), 1.22 ( $m_c$ , 4 H,  $\text{CH}_2$ ), 1.32 (t,  $J = 7.7$  Hz, 1 H, SH), 1.38 (quin,  $J = 7.4$  Hz, 2 H,  $\text{CCH}_2\text{CH}_2$ ), 1.44 ( $m_c$ ,  $J = 2.5$  Hz, 6 H,  $\text{CCH}_2\text{-Ad}$ ), 1.60 (quin,  $J = 7.4$  Hz, 2 H,  $\text{CH}_2\text{CH}_2\text{SH}$ ), 1.64 ( $m_c$ , 6 H,  $\text{CH}_2\text{-Ad}$ ), 1.92 ( $m_c$ , 3 H,  $\text{CH}_{\text{Ad}}$ ), 2.51 (q,  $J = 7.4$  Hz, 2 H,  $\text{CH}_2\text{SH}$ ).


$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta = 22.3$  ( $\text{CH}_2$ ), 24.8 ( $\text{CH}_2$ ), 28.5 ( $\text{CH}_2$ ), 28.9 ( $\text{CH}_{\text{Ad}}$ ), 30.1 ( $\text{CH}_2$ ), 32.3 ( $\text{C}_{\text{Ad}}$ ), 34.2 ( $\text{CH}_2$ ), 37.4 ( $\text{CH}_2$ ), 42.6 ( $\text{CH}_{\text{Ad}}$ ), 44.8 ( $\text{CH}_2$ ).

IR (NaCl):  $\nu(\text{cm}^{-1}) = 2902s, 2846s, 2663w, 1450m, 1347w, 1101w$ .

MS (EI):  $m/z$  (rel int %) = 252 ( $\text{M}^+$ , 3), 135 (100), 93 (11), 79 (11), 67 (6), 41 (6).

Elemental analysis for  $\text{C}_{16}\text{H}_{28}\text{S}$  calc.: C 76.12%, H 11.18%; found: C 76.07%, H 11.08%.

## Synthesis of 11-aminoundecanol



11-Aminoundecanoic acid (21.0 g, 102 mmol) was added slowly at -20 °C to a solution of  $\text{LiAlH}_4$  (5.20 g, 137 mmol) in freshly distilled THF (200 mL) and the reaction mixture was stirred for 6 hours at 67 °C. The solution was cooled to 0 °C, treated with a NaOH solution (10%, 10 mL) and  $\text{H}_2\text{O}$  (20 mL) and stirred for 30 min at 23 °C. The reaction mixture was dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure to provide the desired 11-aminoundecanol (10.3 g) in 53% yield.

m.p.: 66 - 68 °C.

$^1\text{H}$ -NMR (400.1 MHz,  $\text{CD}_2\text{Cl}_2$ , 295 K):  $\delta = 1.27$  ( $s_b$ , 14 H,  $\text{CH}_2$ ), 1.39 (quin,  $J = 7.0$  Hz, 2 H,  $\text{CH}_2\text{CH}_2\text{NH}_2$ ), 1.50 (quin,  $J = 6.5$  Hz, 2 H,  $\text{CH}_2\text{CH}_2\text{OH}$ ), 2.61 (t,  $J = 7.0$  Hz, 2 H,  $\text{CH}_2\text{NH}_2$ ), 3.54 (t,  $J = 6.5$  Hz, 2 H,  $\text{CH}_2\text{OH}$ )  $\Rightarrow$  OH und  $\text{NH}_2$  signals lies under the quintetts between 1.36 - 1.54 Hz.


$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $\text{CD}_2\text{Cl}_2$ , 295 K):  $\delta = 25.9$  ( $\text{CH}_2$ ), 27.0 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 29.6 ( $\text{CH}_2$ ), 29.6 ( $\text{CH}_2$ ), 29.7 ( $\text{CH}_2$ ), 29.7 ( $\text{CH}_2$ ), 33.0 ( $\text{CH}_2\text{CH}_2\text{OH}$ ), 34.1 ( $\text{CH}_2\text{CH}_2\text{NH}_2$ ), 42.3 ( $\text{CH}_2\text{NH}_2$ ), 62.7 ( $\text{CH}_2\text{OH}$ ).

IR (KBr):  $\nu$  (cm<sup>-1</sup>) = 3338m, 3086m<sub>b</sub>, 2923s, 2849s, 1614w, 1472m, 1348m, 1077m, 1031m, 991m.

MS (FAB):  $m/z$  (rel int %) = 188 ([M + H]<sup>+</sup>, 100), 69 (10), 55 (19), 44 (9), 41 (9).

Elemental analysis for C<sub>11</sub>H<sub>25</sub>NO calc.: C 70.53%, H 13.45%, N 7.48%; found: C 70.69%, H 13.43%, N 7.35%.

### Synthesis of *tert*-butyl 11-hydroxyundecylcarbamate



Triethylamine (9.00 mL, 64.7 mmol) and di-*tert*-butyl dicarbonate (14.1 g, 64.6 mmol) were added at 0 °C to a solution of 11-aminoundecanol (12.0 g, 64.1 mmol) in freshly distilled CH<sub>2</sub>Cl<sub>2</sub> (200 mL) and the reaction mixture was stirred for 2 hours at 0 °C and another 8 hours at 23 °C. The resulting solution was washed with a saturated NaCl solution (100 mL) and water (100 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with 30% AcOEt in hexanes, gave 89% yield of *tert*-butyl 11-hydroxyundecylcarbamate (16.3 g).

m.p.: 34 - 36 °C. R<sub>f</sub> = 0.38 (hexanes/AcOEt = 2:1).

<sup>1</sup>H-NMR (400.1 MHz, CDCl<sub>3</sub>, 295 K):  $\delta$  = 1.25 (s<sub>b</sub>, 10 H, CH<sub>2</sub>), 1.30 (m<sub>c</sub>, 4 H, CH<sub>2</sub>), 1.42 (s<sub>b</sub>, 11 H, C(CH<sub>3</sub>)<sub>3</sub> + CH<sub>2</sub>CH<sub>2</sub>NH), 1.54 (m<sub>c</sub>, 3 H, OH + CH<sub>2</sub>CH<sub>2</sub>OH), 3.08 (m<sub>c</sub>,  $J$  = 6.3 Hz, 6.6 Hz, 2 H, CH<sub>2</sub>NH), 3.61 (m<sub>c</sub>,  $J$  = 5.6 Hz, 6.3 Hz, 2 H, CH<sub>2</sub>OH), 4.54 (s<sub>b</sub>, 1 H, NH).


<sup>13</sup>C{<sup>1</sup>H}-NMR (100.6 MHz, CDCl<sub>3</sub>, 295 K):  $\delta$  = 25.8 (CH<sub>2</sub>), 26.8 (CH<sub>2</sub>), 28.5 (C(CH<sub>3</sub>)<sub>3</sub>), 29.3 (CH<sub>2</sub>), 29.5 (CH<sub>2</sub>), 29.5 (CH<sub>2</sub>), 29.6 (CH<sub>2</sub>), 29.6 (CH<sub>2</sub>), 30.1 (CH<sub>2</sub>CH<sub>2</sub>NH), 32.8 (CH<sub>2</sub>CH<sub>2</sub>OH), 40.7 (CH<sub>2</sub>NH), 63.1 (CH<sub>2</sub>OH), 79.1 (C(CH<sub>3</sub>)<sub>3</sub>), 156.1 (CO).

IR (NaCl):  $\nu$  (cm<sup>-1</sup>) = 3353m<sub>b</sub>, 2926s, 2854m, 1693s, 1526m<sub>b</sub>, 1366m, 1275m, 1251m, 1173s.

MS (FAB):  $m/z$  (rel int %) = 288 ([M + H]<sup>+</sup>, 3), 232 (7), 188 (17), 74 (12), 69 (11), 57 (100), 41 (26).

Elemental analysis for C<sub>16</sub>H<sub>33</sub>NO<sub>3</sub> calc.: C 66.86%, H 11.57%, N 4.87%; found: C 66.95%, H 11.49%, N 4.79%.

### Synthesis of *tert*-butyl 11-bromoundecylcarbamate



*N*-Bromosuccinimide (10.0 g, 56.2 mmol) and triphenylphosphine (14.8 g, 56.4 mmol) were added to a solution of *tert*-butyl 11-hydroxyundecylcarbamate (16.0 g, 55.7 mmol) in freshly distilled benzene (200 mL) and the reaction mixture was stirred for 12 hours at 23 °C. The resulting solution was washed with a saturated NaCl solution (200 mL) and water (200 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. Purification by flash chromatography, eluting with 10% AcOEt in hexanes, gave 74% yield of *tert*-butyl 11-bromoundecylcarbamate (14.5 g).

m.p.: 60 - 62 °C.  $R_f$  = 0.37 (hexanes/AcOEt = 10:1).

$^1\text{H}$ -NMR (400.1 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 1.25 ( $s_b$ , 12 H,  $\text{CH}_2$ ), 1.42 ( $s_b$ , 13 H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$  +  $\text{C}(\text{CH}_3)_3$  +  $\text{CH}_2\text{CH}_2\text{NH}$ ), 1.83 ( $m_c$ ,  $J$  = 6.9 Hz, 7.2 Hz, 2 H,  $\text{CH}_2\text{CH}_2\text{Br}$ ), 3.08 (t,  $J$  = 6.6 Hz, 2 H,  $\text{CH}_2\text{NH}$ ), 3.38 (t,  $J$  = 6.9 Hz, 2 H,  $\text{CH}_2\text{Br}$ ), 4.53 ( $s_b$ , 1 H,  $\text{NH}$ ).


$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 26.8 ( $\text{CH}_2$ ), 28.2 ( $\text{CH}_2$ ), 28.5 ( $\text{C}(\text{CH}_3)_3$ ), 28.8 ( $\text{CH}_2$ ), 29.3 ( $\text{CH}_2$ ), 29.4 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 30.1 ( $\text{CH}_2\text{CH}_2\text{NH}$ ), 32.9 ( $\text{CH}_2\text{CH}_2\text{Br}$ ), 34.1 ( $\text{CH}_2\text{Br}$ ), 40.7 ( $\text{CH}_2\text{NH}$ ), 79.0 ( $\text{C}(\text{CH}_3)_3$ ), 156.0 (CO).

IR (KBr):  $\nu$  ( $\text{cm}^{-1}$ ) = 3377 $s$ , 2918 $s$ , 2851 $s$ , 1687 $s_b$ , 1522 $s_b$ , 1469 $m$ , 1374 $m$ , 1280 $m$ , 1237 $m$ , 1170 $m_b$ , 1026 $w$ , 976 $w$ , 870 $m$ , 720 $m$ , 643 $m$ .

MS (FAB):  $m/z$  (rel int %) = 350 ( $\text{M}^+$ , 4,  $^{79}\text{Br}$ ), 294 (96,  $^{79}\text{Br}$ ), 250 (13,  $^{79}\text{Br}$ ), 214 (8), 137 (8), 74 (41), 57 (100), 41 (20).

Elemental analysis for  $\text{C}_{16}\text{H}_{32}\text{BrNO}_2$  calc.: C 54.85%, H 9.21%, N 3.91%; found: C 55.01%, H 9.06%, N 3.89%.

### Synthesis of (*S*)-11-(*tert*-butoxycarbonyl)undecylethanethioate



Thioacetic acid (4.00 mL, 56.0 mmol), triethylamine (8.00 mL, 57.5 mmol) and 4-DMAP (0.50 g, 4.10 mmol) were added to a solution of *tert*-butyl 11-bromoundecylcarbamate (14.5 g, 41.4 mmol) in freshly distilled  $\text{CH}_2\text{Cl}_2$  (200 mL) and the reaction mixture was stirred for 24 hours at 23 °C. The solution was washed with a saturated  $\text{Na}_2\text{CO}_3$  solution (200 mL) and water (200 mL). The organic phase was dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. Purification of the crude material by flash chromatography, eluting with 10% AcOEt in hexanes and subsequent crystallization (*n*-pentane), gave 79% yield of (*S*)-11-(*tert*-butoxycarbonyl)undecyl-ethanethioate (11.3 g).

m.p.: 64 - 66 °C.  $R_f$  = 0.25 (hexanes/AcOEt = 10:1).

$^1\text{H}$ -NMR (400.1 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 1.22 ( $s_b$ , 10 H,  $\text{CH}_2$ ), 1.25 ( $m_c$ , 2 H,  $\text{NCH}_2\text{CH}_2\text{CH}_2$ ), 1.30 ( $m_c$ , 2 H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{S}$ ), 1.41 ( $s_b$ , 11 H,  $\text{C}(\text{CH}_3)_3$  +  $\text{NCH}_2\text{CH}_2$ ), 1.52 ( $m_c$ ,  $J$  = 6.8 Hz, 7.4 Hz, 2 H,  $\text{CH}_2\text{CH}_2\text{S}$ ), 2.29 (s, 3 H,  $\text{SCOCH}_3$ ), 2.83 (t,  $J$  = 7.4 Hz, 2 H,  $\text{CH}_2\text{S}$ ), 3.07 (t,  $J$  = 6.6 Hz, 2 H,  $\text{NCH}_2$ ), 4.55 ( $s_b$ , 1 H,  $\text{NH}$ ).

$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $\text{CDCl}_3$ , 295 K):  $\delta$  = 26.8 ( $\text{NCH}_2\text{CH}_2\text{CH}_2$ ), 28.5 ( $\text{C}(\text{CH}_3)_3$ ), 28.8 ( $\text{CH}_2\text{S}$ ), 29.1 ( $\text{CH}_2$ ), 29.2 ( $\text{CH}_2$ ), 29.3 ( $\text{CH}_2\text{CH}_2\text{S}$ ), 29.4 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 29.5 ( $\text{CH}_2$ ), 30.1 ( $\text{NCH}_2\text{CH}_2$ ), 30.7 ( $\text{SCOCH}_3$ ), 40.6 ( $\text{NCH}_2$ ), 78.9 ( $\text{C}(\text{CH}_3)_3$ ), 156.0 (CO), 196.0 (SCO).


IR (KBr):  $\nu$  ( $\text{cm}^{-1}$ ) = 3380 $m$ , 2918 $m$ , 2851 $m$ , 1687 $s_b$ , 1522 $s_b$ , 1469 $m$ , 1364 $m$ , 1282 $m$ , 1238 $m$ , 1172 $m$ , 1138 $m$ , 1116 $m$ , 1000 $w$ , 959 $m$ , 870 $w$ , 720 $w$ , 639 $m$ .

MS (FAB):  $m/z$  (rel int %) = 346 ( $[\text{M} + \text{H}]^+$ , 24), 290 (26), 246 (100), 230 (9), 202 (9), 57 (55), 43 (31).

Elemental analysis for  $\text{C}_{18}\text{H}_{35}\text{NO}_3\text{S}$  calc.: C 62.56%, H 10.20%, N 4.05%; found: C 62.56%, H 10.16%, N 4.08%.



## Synthesis of *tert*-butyl 11-mercaptoundecylcarbamate



A hydrazine solution (1 M in THF, anhydrous, 30.0 mL, 30.0 mmol) was added at 0 °C to a solution of (*S*)-11-(*tert*-butoxycarbonyl)undecylethanethioate (1.00 g, 2.90 mmol) in freshly distilled CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and the reaction mixture was stirred for 4 hours at 23 °C. The solvent was concentrated under reduced pressure and the residue dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The resulting solution was washed with a saturated NH<sub>4</sub>Cl solution (50 mL) and water (50 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. Purification by flash chromatography, eluting with 10% AcOEt in hexanes, gave 98% yield of *tert*-butyl 11-mercaptoundecylcarbamate (0.86 g).

m.p.: 52 - 54 °C.  $R_f$  = 0.36 (hexanes/AcOEt = 10:1).

<sup>1</sup>H-NMR (400.1 MHz, CDCl<sub>3</sub>, 295 K):  $\delta$  = 1.25 (s<sub>b</sub>, 12 H, CH<sub>2</sub>), 1.32 (t,  $J$  = 7.6 Hz, 1 H, SH), 1.36 (m<sub>c</sub>, 2 H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>SH), 1.43 (s<sub>b</sub>, 11 H, C(CH<sub>3</sub>)<sub>3</sub> + NCH<sub>2</sub>CH<sub>2</sub>), 1.59 (quin,  $J$  = 7.2 Hz, 2 H, CH<sub>2</sub>CH<sub>2</sub>S), 2.51 (m<sub>c</sub>,  $J$  = 7.2 Hz, 7.6 Hz, 2 H, CH<sub>2</sub>SH), 3.09 (t,  $J$  = 6.6 Hz, 2 H, NCH<sub>2</sub>), 4.50 (s<sub>b</sub>, 1 H, NH).


<sup>13</sup>C{<sup>1</sup>H}-NMR (100.6 MHz, CDCl<sub>3</sub>, 295 K):  $\delta$  = 24.7 (CH<sub>2</sub>SH), 26.9 (CH<sub>2</sub>), 28.4 (CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>SH), 28.5 (C(CH<sub>3</sub>)<sub>3</sub>), 29.1 (CH<sub>2</sub>), 29.3 (CH<sub>2</sub>), 29.5 (CH<sub>2</sub>), 29.5 (CH<sub>2</sub>), 29.6 (CH<sub>2</sub>), 30.1 (NCH<sub>2</sub>CH<sub>2</sub>), 34.1 (CH<sub>2</sub>CH<sub>2</sub>SH), 40.7 (NCH<sub>2</sub>), 79.0 (C(CH<sub>3</sub>)<sub>3</sub>), 156.0 (CO).

IR (KBr):  $\nu$  (cm<sup>-1</sup>) = 3373s, 2921s, 2848s, 1682s<sub>b</sub>, 1513s<sub>b</sub>, 1464m, 1367m, 1284m, 1241m, 1166s<sub>b</sub>, 1044w, 1001m, 977m, 864m, 783w, 723m, 580m<sub>b</sub>.

MS (FAB):  $m/z$  (rel int %) = 304 ([M + H]<sup>+</sup>, 10), 248 (41), 204 (100), 74 (13), 57 (77), 41 (14).

Elemental analysis for C<sub>16</sub>H<sub>33</sub>NO<sub>2</sub>S calc.: C 63.32%, H 10.96%, N 4.62%, O 10.54%; found: C 63.41%, H 10.90%, N 4.62%, O 10.46%.

## Synthesis of 11-aminoundecane-1-thiol hydrochloride



*Tert*-butyl 11-mercaptoundecylcarbamate (1.00 g, 3.30 mmol) was dissolved in a HCl solution in dioxane (4 M, 20.0 mL, 80 mmol) and the reaction mixture was stirred for 3 hours at 23 °C. The solvent was concentrated under reduced pressure and the solid washed with diethyl ether (50 mL) and *n*-pentane (50 mL) to provide the desired 11-aminoundecane-1-thiol hydrochloride (0.78 g) in 99% yield.

m.p.: 162 - 164 °C.

<sup>1</sup>H-NMR (400.1 MHz, (CD<sub>3</sub>)<sub>2</sub>SO, 295 K):  $\delta$  = 1.25 (s<sub>b</sub>, 12 H, CH<sub>2</sub>), 1.32 (m<sub>c</sub>, 2 H, NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.51 (quin,  $J$  = 6.8 Hz, 2 H, CH<sub>2</sub>CH<sub>2</sub>SH), 1.53 (quin,  $J$  = 7.4 Hz, 2 H, NCH<sub>2</sub>CH<sub>2</sub>), 2.23 (t,  $J$  = 7.6 Hz, 1 H, SH), 2.46 (m<sub>c</sub>,  $J$  = 6.8 Hz, 7.6 Hz, 2 H, CH<sub>2</sub>SH), 2.75 (t<sub>b</sub>,  $J$  = 7.4 Hz, 2 H, NCH<sub>2</sub>), 7.78 (s<sub>b</sub>, 3 H, NH<sub>3</sub>Cl).

$^{13}\text{C}\{^1\text{H}\}$ -NMR (100.6 MHz,  $(\text{CD}_3)_2\text{SO}$ , 295 K):  $\delta$  = 23.7 ( $\text{CH}_2\text{SH}$ ), 25.8 ( $\text{CH}_2$ ), 26.9 ( $\text{NCH}_2\text{CH}_2$ ), 27.7 ( $\text{NCH}_2\text{CH}_2\text{CH}_2$ ), 28.5 ( $\text{CH}_2$ ), 28.5 ( $\text{CH}_2$ ), 28.8 ( $\text{CH}_2$ ), 28.9 ( $\text{CH}_2$ ), 28.9 ( $\text{CH}_2$ ), 33.4 ( $\text{CH}_2\text{CH}_2\text{SH}$ ), 38.7 ( $\text{NCH}_2$ ).

IR (KBr):  $\nu$  ( $\text{cm}^{-1}$ ) = 2918<sub>s</sub>, 2657<sub>m</sub>, 2547<sub>m</sub>, 2021<sub>w</sub>, 1625<sub>m</sub>, 1511<sub>m</sub>, 1473<sub>m</sub>, 1398<sub>w</sub>, 1308<sub>w</sub>, 1196<sub>w</sub>.


MS (FAB):  $m/z$  (rel int %) = 204 ( $[\text{M} - \text{Chlorid}]^+$ , 100), 55 (6).

Elemental analysis for  $\text{C}_{11}\text{H}_{26}\text{ClNS}$  calc.: C 55.09%, H 10.93%, N 5.84%; found: C 55.21%, H 10.89%, N 5.73%.

### Analyses of thiolate-protected gold colloids

Colloid	d (TEM) [nm]	Elemental analysis
<b>6</b>	$2.32 \pm 0.46$ nm	C 16.52%, S 2.29%, Au 79.5%
<b>7</b>	$3.48 \pm 0.61$ nm	C 7.15%, S 2.38%, Au 90.1%
<b>8</b>	$3.44 \pm 0.59$ nm	C 11.39%, S 2.22%, Au 76.8%
<b>9</b>	$2.60 \pm 0.43$ nm	C 7.39%, N 0.71, S 1.40%, Au 81.5%
<b>10</b>	$3.39 \pm 0.73$ nm	C 8.26%, O 0.83, S 1.79%, Au 79.3%

TEM-image of colloid 7



### Complete Reference 13:

- [13] M. J. Hostetler, J. E. Wingate, C.-J. Zhong, J. E. Harris, R. W. Vachet, M. R. Clark, J. D. Londono, S. J. Green, J. J. Stokes, G. D. Wignall, G. L. Glish, M. D. Porter, N. D. Evans, R. W. Murray, *Langmuir* **1998**, *14*, 17.